


Welcome to the Second Annual General Meeting Wetlands as Nature-Based Climate Solutions

May 1, 2024

# Welcome



# **CAAF Project:**

Wetlands as nature-based climate-change solutions: Quantifying carbon-capture potential while building a stronger green economy.



# **CAAF**, All Objectives

| 2023 | 2024 | 2025 | 2026 | 2027

Start April 1, 2022

**Core Activities** 

April 1, 2022 - March 31, 2027

Finish March 31, 2027

Objective 1. Develop Authoritative Estimates Of Landscape-Scale Density Of Wetland Coverage For Agricultural Landscapes

April 1, 2022 - March 31, 2025

Objective 2. Develop Authoritative Estimates For Rates Of OC Accumulation, GHG Fluxes To The Atmosphere, And Carbon Transports Into (And Out Of) Wetlands

April 1, 2022 - October 15, 2026

Objective 3. Develop Robust Estimates Of Hydrological Process Controls On OC Accumulation And GHG Fluxes From Wetlands

April 1, 2022 - March 31, 2027

Objective 4. Develop Robust Estimates Of The Synergies (And Conflicts) Of Wetlands As NBS For Carbon Storage Versus Other Benefits

April 1, 2022 - March 31, 2027

Objective 5. Use The Authoritative And Robust Estimates Of OC Accumulation And GHG Fluxes To Inform Policy And Practice Tools To Incentivize The Use Of Wetlands As NBS For Multiple Benefits In Agricultural Landscapes

April 1, 2023 - March 31, 2027

# **Prairies Project:**

Wetland as Natural Solutions: Baselines and projections for Wetlands on Agricultural Land



# **Prairies**

**Finish** 2025 2024 2023 March 31, 2025 Start April 1, 2022 Objective 1. Identify the main drivers of wetland conversion and projecting how these drivers might change the rates of wetland conversion, and the resulting GHG emissions, over time. Objective 2. Analysis of how activities might shift to a different location or ecosystem (leakage) if activities are undertaken to reduce the draining of wetlands on agricultural lands in each region. Task 1. Conceptual socioeconomic model planning approach for Lake Winnipeg watershed Task 2. Engagement with science teams in the development of wetland inventory Task 3. Analysis of primary drivers of wetland conversion Task 4. Development of survey instrument on socioeconomic perspectives Task 5. Development of socioeconomic model for wetland conversion Task 6. Evaluation and assessment of policy instruments Task 7. Development of model to assess leakage of wetlands Task 8. Research Team

## **Great Lakes Project**

Wetlands as Nature-Based Climate Solutions: A Socioeconomic Analysis of the Great Lakes-St. Lawrence River Basin



# Great Lakes, all objectives

Start

| Jan '24 | Apr '24 | Jul '24 | Oct '24 | Jan '25 | Apr '25 | Jul '25 | Oct '25 | Jan '26 | Apr '26 | Jul '26 | Oct '26 |Jan '27

Finish January 4, 2027

January 1, 2024 Wetlands as Nature-Based Climate Solutions: A Socioeconomic Analysis of the **Great Lakes-St. Lawrence River Basin** January 1, 2024 - January 4, 2027


> Objective 1. Evaluate methods of mapping wetlands and estimating historical rates of land use change and conversion of wetlands in the GLSLRB January 1, 2024 - December 31, 2024

> > Objective 2. Create an inventory of wetlands and identify historical rates of land use change and conversion of wetlands in the GLSLRB January 1, 2025 - January 1, 2027

Objective 3. Identify the main socio-economic drivers of wetland conversion and projecting how these drivers might change the rates of wetland conversion, and the resulting GHG emissions, over time in the GLSLRB January 1, 2024 - January 4, 2027

Objective 4. Cost-effectiveness and cost-benefit analysis of restoration and/or conservation of wetlands on agricultural lands as NbCS in the GLSLRB January 1, 2024 - January 4, 2027

# National Wetland Information Network



# Investigators (CAAF Project)



# Investigators (Prairies Project)



Dr. Irena Creed



Dr. Patrick Lloyd-Smith



**Dr. John Pattison-Williams** 

# Investigators (Great Lakes Project)







Dr. Georgios Arhonditsis



**Dr. Roy Brouwer** 



Dr. Jie He



Dr. Ben DeVries



Dr. Geneviève Ali



Dr. Lota Tamini

# Wetland Assessments

Area covered by: CAAF Wetlands as Nature Based Climate Solutions (green)

Lake Winnipeg Watershed Area covered by the first extension to the

CAAF Wetlands as Nature Based Climate Solutions Project

led by Patrick Lloyd-Smith, and John Pattison-Williams.

Great Lakes-St. Lawrence River Basin area\* covered by the proposed second extension to the CAAF Wetlands as Nature Based Climate Solutions Project

led by
Roy Brouwer
Jie He
Ben DeVries
Genèvieve Ali
George Arthonditsis
Lota Tamini

Source: Statistics Canada, Agriculture Division, Remote Sensing and Geospatial Analysis section, 2017. Agricultural Ecumene Boundary File – 2016.

https://open.canada.ca/data/en/dataset/317bf695-b6e2-4b60-90a8-51cd3c3d3d64



Area covered by: CAAF Wetlands as Nature Based Climate Solutions (green)

Lake Winnipeg Watershed Area covered by the first extension to the

CAAF Wetlands as Nature Based Climate Solutions Project

led by Patrick Lloyd-Smith,

and John Pattison-Williams.

Cross fertilization of theories, methods, approaches

Great Lakes-St. Lawrence River Basin area\* covered by the proposed second extension to the CAAF Wetlands as Nature Based Climate Solutions Project led by

> Roy Brouwer Jie He Ben DeVries Genèvieve Ali

George Arthonditsis

14

Source: Statistics Canada, Agriculture Division, Remote Sensing and Geospatial Analysis section, 2017. Agricultural Ecumene Boundary File – 2016.

https://open.canada.ca/data/en/dataset/317bf695-b6e2-4b60-90a8-51cd3c3d3d64

# Agenda, May 1

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- · 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks



# Agenda, May 1

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- 3:30 − 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

### Rose Fuchsia Room, Topic A

### Purple Room, Topic B

**Topics:** 

GIS, remote sensing, mapping, and modeling

**Objectives:** 

CAAF, Objectives 1, 3, 5.3.

Great Lakes, Objectives 1, 2, 4.5.

**Room leaders:** 

David Aldred, Ben DeVries

**Topics:** 

Flux towers, measurements, and carbon cycling

**Objectives:** 

CAAF, Objective 2

**Room leaders:** 

Pascal Badiou, Matthew Bogard

### Rose Fuchsia Room, Topic A

### **Topics:**

GIS, remote sensing, mapping, and modeling

### **CAAF:**

Ali Ameli

George Arhonditsis (Ratnajit Saha, Yoji Uno, Xin Wey Wen)

Irena Creed (David Aldred, Forough Fendereski, Eric Enanga, Shizhou Ma, Nayyer Mirnasl, Michael Dallosch, Pengfei Ren, Adan Auyeung)

### **Great Lakes:**

Ben DeVries (Maciej Lizak, Alejandro Nieto, Kathleen B.)

Genevieve Ali

### Purple Room, Topic B

### **Topics:**

Flux towers, measurements, and carbon cycling

### **CAAF:**

Pascal Badiou

Christian von Sperber (Calder Jones, Shir Gruber)

Matthew Bogard (Chin Ngai Chan)

Larry Flanagan\* (Oshini Fernando)

David Lobb

Gail Chmura (Wendy Ampuero, Rachel Plant)

Sara Knox

# Agenda, May 1

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- · 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

# Agenda, May 1

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

### Room A. Rose Fuchsia

### **Topics:**

GIS, remote sensing, mapping, and modeling

### Room B. Purple

### **Topics:**

Flux towers, measurements, and carbon cycling

### **Ouestions:**

How do we link our measurements and modelling approaches across scales? (i.e. results from soil/water incubations - Kayak sampling - Eddy co-variance measurements on site and across country - large scale modelling approaches).

What information/knowledge do we gain and/or lose during the upscaling? What gaps of knowledge need to be filled to link the different scales?

Discussions on sampling GHG concentrations in water.

# Agenda, May 1

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break Checkout for some travelers
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

# 12:00 – 1:00, Lunch break

Foyer 2
Checkout for some travelers



Welcome to the Second Annual General Meeting Wetlands as Nature-Based Climate Solutions

May 1, 2024



## **ROOM A: Purple Room**

GIS, remote sensing, mapping, and modeling

# ROOM A: GIS, remote sensing, mapping, and modeling

- . Sheel Bansal
- David Aldred
- · Forough Fendereski
- · Eric Enanga
- · Shizhou Ma
- · Nayyer Mirnasl
- · Ratnajit Saha
- Ben DeVries
- . Genevieve Ali
- . David Lobb

# ROOM A: GIS, remote sensing, mapping, and modeling · Sheel Bansal

# Wetland Biogeochemistry: Mechanisms, Models, Management, Methods (Web4M)

U.S. Geological Survey, Climate Research & Development Program

Northern Prairie Wildlife Research Center Jamestown, North Dakota, USA


Sheel Bansal, Senior scientist Brian Tangen, Junior scientist Jacob Meier, Lead technician Max Post van der Burg, Modeler

### 1. Mechanisms



Vegetation, hydrology, and soils

### 2. Models



Spatially explicit methane model

### 3. Management



**Conservation Reserve Program** 

### 4. Methods




Automation, Patents, Student led

ROOM A: GIS, remote sensing, mapping, and modeling · David Aldred

# **Objective 1, Task 2:** Develop authoritative estimates of landscapes scale density of wetland coverage for agricultural landscapes Irena Creed, David Aldred


In Task 1 we compiled existing inventories in the Prairies and identified limitations of having non-standardized approaches (different source data, different dates, different specifications)

In Task 2 we are creating standardized inventories that span multiple years




We tested multi-year (1 to 10 year) overlays of Landsat-based Dynamic Surface Water of Canada\* annual inundation maps with lakes and rivers removed for spatial accuracy and drainage loss estimates in 30 areas where high resolution DUC inventories were available for comparison

Our testing resulted in 28-year (1993-2020) time series of annual wetland inventories based on 10-year overlays




\*Olthof I, Rainville T. 2022. Dynamic surface water maps of Canada from 1984 to 2019 Landsat satellite imagery. Remote Sensing of Environment 279: 113121. https://doi.org/10.1016/j.rse.2022.113121 We are continuing to test different multi-year overlays using frequency of inundation thresholds

Together with Objective 2 of the Great Lakes-St. Lawrence River Basin extension project, we will also test sub-pixel water fraction (SWF) methods\* to improve wetland mapping




\*DeVries B, Huang C, Lang MW, Jones JW, Huang W, Creed IF, Carroll ML. 2017. Automated quantification of surface water inundation in wetlands using optical satellite imagery. Remote Sensing 9(8): 807. https://doi.org/10.3390/rs9080807


Using these inventories and AAFC land cover maps, we generated estimates of number and area of wetland conversions between 1970-1990, 1990-2005, and 2005-2020




Standardized wetland inventories will be available to all project investigators and will directly support Tasks 1.3, 1.4, 3.2, 3.3, 4.2, and 4.4 of the CAAF project and Objectives 1 and 2 of the Prairies Project, and the methods will support Objectives 1 and 2 in the Great Lakes Project



# ROOM A: GIS, remote sensing, mapping, and modeling · Forough Fendereski




We identified wetland "objects" from Landsat inundated areas and extracted wetlandscape properties —number, size, and wetland-to-wetland connectivity— to track their changes from 1984-2020



Fendereski F, Ma S, Mohammady S, Spence C, Trick C, Creed I. Tracking changes in wetlandscape properties of the Lake Winnipeg Watershed using Landsat inundation products (1984-2020). Manuscript in preparation.

Wetlandscape properties increased 1984-2020 in response to increases in precipitation



**Spearman's rank correlation** showed precipitation alone could explain **more than 70%** of the variations in wetlandscape properties (p < 0.01).

# ROOM A: GIS, remote sensing, mapping, and modeling · Eric Enanga

**OBJECTIVE 2.** Develop Authoritative Estimates For Rates Of <u>OC Accumulation</u>, GHG Fluxes To The Atmosphere, And Carbon Transports Into (And Out Of) Wetlands.

**2.2** Develop Standards And Protocols To Measure Wetland <u>OC Accumulation</u> And GHG Flux Rates.

# OC stock estimation: Does the method matter?

CAAF Annual General Meeting, May 1st 2024

Eric Enanga


Supervisor – Dr. Irena Creed

### Equivalent mass vs Depth and <sup>137</sup>Cs 1963 peak OC stock


Method	Depth	Year	Mass
Depth based	Fixed	Variable	Variable
<sup>137</sup> Cs 1963 peak based	Variable	Fixed	Variable
<b>Equivalent Mass based</b>	Variable	Variable	Fixed


Before compaction


After compaction "e.g., tractor tire"



# Depth, <sup>137</sup>Cs 1963 peak & Equivalent mass OC stock estimates







### Take home message

All three methods have a role to play in OC stock estimates:

- Depth based method is fast and efficient where the sampling sites are homogenous with minimal compaction, excellent for screening large sample sets.
- \* <sup>137</sup>Cs 1963 peak method is useful where, in addition, OC sequestration rates are important (i.e., time marker).
- Equivalent mass is much more reliable where substantial compaction is anticipated, e.g., drained sites, although requires more time investment.

### Contributions to Objective 2

- 2.2 Develop Standards And Protocols To Measure Wetland <u>OC</u>

  <u>Accumulation</u> And GHG Flux Rates (April 3, 2023 April 1, 2024).

  <u>We have demonstrated that the Equivalent Mass method is more reliable for consideration as a potential protocol for OC accumulation estimation.</u>
- 2.3 Using Standards And Methodologies Developed In Task 2.2, Measure Wetland *OC Accumulation* And GHG Flux Rates (April 1, 2022 October 15, 2026).


Where substantial compaction is anticipated, equivalent mass should, in future, be the preferred protocol for calculating OC accumulation based on the conclusions arrived at in 2.2 above.


<sup>1.</sup> Creed, I.F., Badiou, P., Enanga, E., Lobb, D.A., Pattison-Williams, J.K., Lloyd-Smith, P., Gloutney, M. (2022). Can restoration of freshwater mineral soil wetlands deliver nature-based climate solutions to agricultural landscapes? *Frontiers in Ecology and Evolution*, 10, 932415, doi:10.3389/fevo.2022.932415, 2022.

<sup>2.</sup> Purbasha, M., Creed, I.F., Trick, C.G., Enanga, E., Lobb, D.A., Submitted. Technical Note: Comparison of radiometric techniques for estimating recent organic carbon sequestration rates in freshwater mineral soil wetlands 3. Ellert B.H., and Bettany J.R. (1995). Calculation of organic matter and nutrients stored in soil under contrasting management regimes. *Canadian Journal of Soil Science* 75: 529-538, doi:10.4141/cjss95-075

# **END** THANK YOU

### Trends in OC stock – Polynomial Orthogonal Contrasts






ROOM A: GIS, remote sensing, mapping, and modeling · Shizhou Ma

### Objective 2: Develop authoritative estimates for rates of organic carbon accumulation, greenhouse gas fluxes to the atmosphere, and carbon transports into (and out of) wetlands

Task 2.5: Develop models to predict the potential for wetlands for OC sequestration and GHG reduction




Ma, S., Mistry, P., Badiou, P., Bansal, S., Creed, I.F. Factors regulating the potential for freshwater mineral soil wetlands to function as natural climate solutions. Wetlands. (submitted)


### GIS/remote sensing proxies for proximal controls (inspired from the conceptual diagram)


Proximal controls	GIS/remote sensing proxies		
Carbon substrate quantity -	Normalized Difference Vegetation Index (NDVI) and Wetland area		
Carbon substrate quality -	→ Normalized Difference Vegetation Index (NDVI)		
Temperature -	Temperature - Landsat Thermal Bands		
Soil microbial community -	Agricultural intensity – Upland Type (cropland vs grassland)		
Plant community -	Normalized Difference Vegetation Index (NDVI)		
Cation exchange capacity	Cation Exchange Capacity (CEC)		
Aggregate reactivity -	Bulk Density (BD)		
Redox potential -			

### Response variable for the random forest model - wetland carbon sequestration rate:

- 210 Pb Constant Flux (CF) model derived C sequestration rate
- Varying C sequestration rate available for single wetland at different time intervals
- Total 89 observations across Alberta, Saskatchewan and Manitoba








### Random forest partial dependence plot for predictors:


- Functional relationship between predictors and carbon sequestration rate match findings in our review paper



### Next steps

- Find better land use data for North America to upgrade "upland" predictors
- Apply the process informed RF model to understand how different environmental factors affect wetland GHG fluxes
- Predict and upscale carbon sequestration rate and GHG fluxes to the entire PPR







# ROOM A: GIS, remote sensing, mapping, and modeling · Nayyer Mirnasl




# CAAF Objectives



Obj2, Task 2.4: Estimate Lateral Flows of Carbon into (and out of) Wetlands



Obj3, Task 3.1: Develop Mechanistic Models of Carbon Cycling



### **Objective**

Estimating the rates of dissolved organic and inorganic carbon accumulation and transport to and from wetlands




Document spatial and temporal variations in DOC and DIC concentrations in the stream network



Identify causes of such variability



Gain insights into the relationship between wetlands and DOC/DIC export from the basin



### **Study** area




Canadian side of the Lake Erie **Basin** 



10.5% increase in downstream DOC export relative to terrestrial inputs



DIC export shows 0.5% increase relative to terrestrial inputs



(Xenopoulos, 2017)

### **SPARROW (SPAtially Referenced** Regressions On Watershed attributes)



A parsimonious hybrid empirical/processbased model



W Developed by the United States Geological Survey (USGS)




Suitable for large spatial scales



Rank contributing catchments based on the loads and yields





### **Next Steps**

Estimating the hydrological connectivity of wetlands and assessing its influence on DIC/DOC fate

Modeling the carbon cycle in wetlands to comprehend the behavior of various wetland classes

Determining wetlands as either sinks or sources of carbon

# ROOM A: GIS, remote sensing, mapping, and modeling · Ratnajit Saha



# Quantify agricultural impact on GHG emissions at major cash-crop farms in a changing climate in Ontario, Canada

Supervisor: Prof. George B. Arhonditsis

#### Ratnajit Saha

PhD Student Ecological Modelling Laboratory Department of Physical & Environmental Sciences University of Toronto Scarborough, Toronto, Ontario, Canada



agriculture.canada.ca

### **Objective**


Aims to investigate the impact of farming activities on GHG emissions in a changing climate in Ontario over the past two decades

### **Methodology-Data sources**

- Crop rotation practices from Statistics Canada and OMAFRA databases
- Identifying agriculture-intensive counties across Ontario
- Novel concept of virtual farm is used (for a first time) to quantify GHG emissions

### **Significance**

- Provincial and national agricultural strategy to achieve "net zero-carbon emission".
- ➤ Develop nature-based solutions to support carbon-smart food production systems and be a part of the national strategy on agri-food sustainability.



An example of a virtual farm


### **Agriculture-Intensive Counties**

#### Crop intensive

- 1. Huron
- 2. Chatham-Kent
- 3. Middlesex
- 4. Lambton
- 5. Perth

#### Livestock intensive

- 1. Wellington
- 2. Perth
- 3. Waterloo
- 4. Bruce
- 5. Kawartha Lake




### **Holos model**

- ☐ Model based on empirical flux estimates
- □ Primarily based on IPCC Tier 2 (2006) methodology: modified to reflect Canadian conditions
- ☐ Yearly and seasonal time step
- ☐ Farm-level scale
- ☐ Boundaries of the system are the farm gate

#### **Uses of Holos model**

- Understand, predict and control a food-production system
- ➤ Identify areas of deficient knowledge
- ➤ Answer various "what if?" scenarios
- ➤ Adding value to experiments

To estimate GHG emissions from farms and test best management practices to reduce emissions



agriculture.canada.ca




https://agriculture.canada.ca/en/agricultural-production/holos-software-program


### **County-wise total GHG emissions**

GHG fluxes from agricultural virtual farms

Years: 2001, 2006, 2011 and 2016


❖ Total-farm emission: Crops versus livestock






Year: 2001

(Indirect  $N_2O$ > Enteric  $CH_4$ > Direct  $N_2O$  > Manure  $CH_4$  > Farm energy  $CO_2$ )




Year: 2006

(Indirect  $N_2O$ > Enteric  $CH_4$ > Direct  $N_2O$  > Manure  $CH_4$  > Farm energy  $CO_2$ )




Year: 2011

(Indirect  $N_2O$ > Enteric  $CH_4$ > Direct  $N_2O$  > Manure  $CH_4$  > Farm energy  $CO_2$ )



Year: 2016


(Indirect  $N_2O$ > Enteric  $CH_4$ > Direct  $N_2O$  > Manure  $CH_4$  > Farm energy  $CO_2$ )



#### **Next Steps**

(Lake Erie Basin)


- ☐ Quantify GHG emissions in the rest of the counties.
- Explore the role of wetlands for carbon sequestration.
- □ Validate GHG fluxes derived from the HOLOS model.



#### **Next Steps**

(Lake Erie watershed area)

- ☐ Quantify GHG emissions in the rest of the counties.
- Explore the role of wetlands for carbon sequestration.
- □ Validate GHG fluxes derived from the HOLOS model.



## You


#### Get in touch!

Email:

ratnajit.saha@mail.utoronto.ca










Linked-in: www.linkedin.com/in/ratnajit-saha-069922a0

Google Scholar:

https://scholar.google.ca/citations?hl=en&tzom=300&user=kQEPsAEAAAAJ&view\_op=list\_works&sortby=pubdate



# ROOM A: GIS, remote sensing, mapping, and modeling . Ben DeVries

## Quantifying and Characterizing wetland change in the Great Lakes St. Lawrence Basin

Ben DeVries, Maciej Lizak, Alejandro Nieto Rodriguez, Kathleen Reid

Department of Geography, Environment and Geomatics
University of Guelph

ECCC-NbS Annual General Meeting May 1, 2024, Toronto, ON





#### **Project Objectives**


Evaluate satellite-based methods of mapping wetlands and estimating historical rates (1970 to present) of land use change and conversion of wetlands in the Great Lakes-St. Lawrence River Basin.

collaboration with UGuelph & McGill U

- 1. Compile ground-based wetland inventories from governmental or non-governmental sources (e.g., DUC SOLRIS inventory) and reported estimates of uncertainty for these wetland inventories (i.e., minimum mapping size, spatial accuracy, omission/commission errors associated with wetland coverage Lead: University of Toronto, in methodology, unaccounted wetlands (i.e., those too small to be captured)) are identified.
  - 2. Compare ground-based with satellite-based wetland inventories to determine the best available wetland mapping methods and to produce a report (see adjacent column) providing the limitations and uncertainties associated with both methods.
  - 3. Estimate wetland and landcover change from 1970 to present.

Create an inventory of wetlands and identify historical rates (1970 Lead: University of Toronto, in stratified random sample of change types (including strata of no significant change) using very highto present) of land use change and conversion of wetlands in the collaboration with UGuelph & resolution reference imagery. By performing this analysis on a stratified random sample, unbiased areal Great Lakes-St. Lawrence River Basin. McGill U

- 1. Wetlands are mapped using methods tested in Strategy 1. Estimates are provided of historical rates of conversion of wetlands in the Great Lakes-St. Lawrence River Basin. Wetland loss (or gain) is mapped by applying a number of statistical analyses, including non-parametric Theil-Sen/Mann-Kendall trend tests, to each pixel in the SWF image stack. These maps and rates will provide inputs into the next objectives (strategies).
- 2. Where wetlands are lost, its follow-on land use and land cover will be classified using pixels drawn from a estimate of follow-on land uses will be produced, which will inform further analysis of socio-economic drivers of wetland change.
- 3. Estimate changes from 1970 to present in wetland number, size, permanence, perimeter: area ratio, and perimeter width and distinctness (factors known to influence carbon cycling) are provided at several assessment unit levels in the Great Lakes-St. Lawrence River Basin, including the Soil Landscapes of Canada (SLC) soil polygon database, which is currently being used to support the national greenhouse gas (GHG) inventory. Uncertainties associated with land cover change/wetland conversion rate are estimated.





#### Wetlands mapping and inventories




Figure: Maciej Lizak


Identify and evaluate existing datasets:


- DUC inventories
- ECCC-CWI
- Other LULC products


Derive high-confidence cal-val data for wetland (type) classification

Develop automated classification models using combinations of optical and (multi-frequency) SAR data:

- Landsat + Sentinel-2
- Sentinel-1 + ALOS-PALSAR (+ RCM + NISAR)
- Terrain indices











Figure: Kathleen Reid

DeVries et al. (2017)

Genevieve Ali, Ben DeVries, Wanhong Yang (OMAFRA)



#### Time Series decomposition



#### **Hypothesis:**

Abrupt, semi-permanent changes in wetland SWF are likely due to LULC changes

ROOM A: GIS, remote sensing, mapping, and modeling . Genevieve Ali

## Refining the definition and classification of mineral wetlands

#### **Genevieve Ali**


Department of Earth & Planetary Sciences | Department of Geography McGill University

## CAAF project focused on the Great Lakes-St. Lawrence River Basin Research objectives related to our work

Objective	Activities / Tasks
Objective 1. Evaluate satellite-based methods of mapping wetlands and estimating historical rates (1970 to present) of land use change and conversion of wetlands in the Great Lakes-St. Lawrence River Basin.	<b>Task 1.1.</b> Compile ground-based wetland inventories from governmental or non-governmental sources (e.g., DUC SOLRIS inventory) and reported estimates of uncertainty for these wetland inventories (i.e., minimum mapping size, spatial accuracy, omission/commission errors associated with wetland coverage methodology, unaccounted wetlands (i.e., those too small to be captured)
	<b>Task 1.2.</b> Compare ground-based with satellite-based wetland inventories to determine the best available wetland mapping methods
	Task 1.3. Estimate wetland and landcover change from 1970 to present.
Objective 2. Create an inventory of wetlands and identify historical rates (1970 to present) of land use change and conversion of wetlands in the Great Lakes-St. Lawrence River Basin	<b>Task 2.1.</b> Provide estimates are provided of historical rates of conversion of wetlands in the Great Lakes-St. Lawrence River Basin; quantify wetland loss (or gain) by applying a number of statistical analyses, including non-parametric Theil-Sen/Mann-Kendall trend tests, to each pixel in the SWF image stack
	<b>Task 2.2.</b> Where wetlands are lost, identify follow-on land use and land cover using pixels drawn from a stratified random sample of change types (including strata of no significant change) using very high-resolution reference imagery.
	<b>Task 2.3.</b> Estimate changes from 1970 to present in wetland number, size, permanence, perimeter: area ratio, and perimeter width and distinctness (factors known to influence carbon cycling) at several assessment unit levels in the Great Lakes-St. Lawrence River Basin, including the Soil Landscapes of Canada (SLC) soil polygon database, which is currently being used to support the national greenhouse gas (GHG) inventory.

## Satellite-based wetland inventories & best mapping methods Considering additional ancillary data in support of SWF data

### U. Of Guelph team (Devries et al.)





Google Earth (April 2020)

#### Data sources

- Stack of Landsat images
- Ancillary data (Digital Elevation Model, Land use and Land cover map)

#### Our planned contribution

- Consider hydric/drainage status of underlying soil
   Goal(s):
  - ✓ Assess the likelihood of wetland presence based on location-specific soil characteristics
- Match SWF map dates with gridded daily climate data (precipitation, antecedent precipitation, potential evapotranspiration, surface soil moisture)
  - Goal(s):
    - ✓ Differentiate wetlands from swales and nonwetland inundated areas
    - ✓ Differentiate surface-water-dominated from groundwater-dominated wetlands

## Changes from 1970 to present in wetland characteristics Adding to the list of wetland characteristics

"Traditional" wetland characteristics used to monitor historical change: wetland number, size, permanence, perimeter: area ratio

#### Potential, additional wetland characteristics:

- Wetland permanence standardized by climate characteristics
  - Goal(s): differentiating permanent (or quasi-permanent)
     open water in dry versus wet years
- Wetland-to-wetland and wetland-to-stream connectivity
  - Goal(s): quantifying historical changes in the frequency with which surface water flow paths may have facilitate connectivity between a wetland and nearby water bodies




#### Work done to date

#### January 2024 to now (Postdoc Sarah Ariano, Genevieve Ali)

Unified North American Soil Map (Liu et al., 2013)

Variables: topsoil and subsoil data from STATSGO/SSURGO and SLC

The First Dominant Soil Component Area Percentage




Computations underway for soil hydraulic parameters using pedotransfer functions (Saxton & Rawls, 2006)


Gridded climate dataset Daymet
Daily climate data, 1-km resolution,
since 1980

Variables: day length, precipitation, shortwave radiation, SWE, max and max air temperature, water vapor pressure

North America Soil Moisture Dataset Derived from Time-Specific Adaptable Machine Learning Models

Variable: near-surface soil moisture (0-5 cm), 250-m resolution de 250 m, biweekly between 2002 and 2020





ROOM A: GIS, remote sensing, mapping, and modeling . David Lobb

## **Lateral Transfers of soil and sediment into wetlands**











Fig. 3. 9. Original unsmoothed 1-m DEM of one wetland catchment: (a) three-dimensional (3D) view of the catchment, (b) colour relief view of the catchment and (c) geometrical characteristics of the sedimentary features for the shown transect in the colour relief view.





#### **ROOM B:**

Flux Towers, measurements, and carbon cycling

#### **ROOM B:** Flux Towers, measurements, and carbon cycling

- · Pascal Badiou
- Larry Flanagan (Matthew Bogard)
- · Matthew Bogard
- · Sara Knox
- · Christian Von Sperber
- . Gail Chmura

**ROOM B:** Flux Towers, measurements, and carbon cycling · Pascal Badiou

## **OBJECTIVE 2** Develop authoritative estimates for rates of *organic* carbon accumulation, greenhouse gas fluxes to the atmosphere, and carbon transports to (and out of) wetlands to downstream water



Overview of monitoring sites and methodologies



## Monitoring GHG emission from freshwater mineral wetland in agricultural landscapes


Prior to CAAF – 48 sites monitored across PPR (black soil zone / Parklands region) including first two freshwater mineral wetland Eddy Covariance flux tower sites in the PPR

Through the CAAF we have expanded monitoring to more than 100 wetland sites including:

- Brown and Dark Brown soil zone in the PPR (both in AB and SK), expanding to additional sites in southern AB (2024)
- Small restored wetlands in southern Ontario (n = 16)
- DUC project wetlands in the BC interior (n = 16)
- Deployed 3rd flux tower at Oak Hammock Marsh
- Collected cores from 2 MB flux tower sites
- Continuing cattail chamber measurements in 2024
- Adding ebullition measurements at flux tower sites in 2024



### **MB Wetland Flux Tower Sites**










## Manitoba We tland Flux Tower Sites





### **Cattail Chambers & Floating Chambers**










#### Extensive dissolved gas sampling - Prairies

- Surface water dissolved gas
- Ambient air
- In situ water quality (pH, SpCond, temp, DO%)
- Surface water area
- Wind speed and atmospheric pressure
- Water chemistry (sulfate, alkalinity, nutrients)






#### **Biodiversity Monitoring - Prairies**

- •We collected data with ARUs at all black soil zone wetland sites (n = 48) in 2022 and a subsample of sites (n = 16) in brown and dark brown soil zones of AB/SK in 2023.
- •We can estimate biodiversity indices (species richness/alpha diversity, evenness, similar) of birds and amphibians at each site to compare to other ecosystem service metrics.
- •Expanding to additional sites spanning a gradient of production where ARUs are deployed and invertebrate sampling and eDNA monitoring is occurring





#### **CAAF – Southwestern Ontario (I)**



Ontario Small Wetland -Water Quality Study (ha)

Nature Force -Water Quantity Study (ha)

n = 8

n = 8

0.34 ± 0.06


0.19 ± 0.04

 $0.27 \pm 0.04$ 






#### **CAAF** – British Columbia





# **ROOM B:** Flux Towers, measurements, and carbon cycling · Larry Flanagan (Matthew Bogard)






Flanagan et al. (2022) Wetlands




#### Apply Remote Sensing Measurements for Scaling and Historical Perspectives of Fluxes



	Stirling Lake	Frank Lake
Dominant Emergent Macrophyte	Cattail	Bulrush
Aboveground Biomass (g m <sup>-2</sup> )	1560 ± 517	352 ± 55
Leaf Area Index (m <sup>2</sup> m <sup>-2</sup> )	5.3 ± 1.6	2.2 ± 0.9

**ROOM B:** Flux Towers, measurements, and carbon cycling · Matthew Bogard







- Deployed 2 bubble traps in 6 wetlands (2023)
- Vol. of gas in bubble traps measured
- Concentration of CH<sub>4</sub>
   in fresh bubbles
   determined by GC

Higher aquatic productivity does NOT result in higher CH<sub>4</sub> flux.

#### 2. Scaling emergent veg C stocks and links to emissions Sam Woodman, PDF


- Used RGB drone imagery, K-mean clustering to quantify emergent veg.
- Estimated emergent veg. biomass by correlating areal biomass & spectral indices
- Diffusive GHG emissions (CO<sub>2</sub>/CH<sub>4</sub>/N<sub>2</sub>O) from aquatic habitats.





- 1. No link b/w veg and  $CO_2/N_2O$  flux.
- 2. Clear positive link b/w veg and CH<sub>4</sub> flux.

#### 3. In kind project: NECB at Frank Lake, AB: Laura Logozzo, PDF



 Scaling C fluxes to entire wetland complex using aquatic emissions/lat. Flux data + biomass sampling + EC + remote sensing

Sustained aquatic emissions dominate NECB due to effluent processing... A tradeoff in services!

**ROOM B:** Flux Towers, measurements, and carbon cycling · Sara Knox


## PROJECT UPDATE



2024 AGM

Sara Knox\*, Pascal Badiou, Lauren Bortolotti, Nick Lee, Darian Ng, Joyson Ahongshangbam, Zoran Nesic

\*Assistant Professor, Department of Geography, McGill University







NEW WETLAND SITE IN QUEBEC (FALL 2024 ORSPRING 2025)








#### **NEW SITE COMING SOON & POTENTIAL SYNERGIES**




















## COLLABORATIONS WITH DUC



- Isolated grassland marsh
- Dominated by emergent vegetation (Tule)
- Characterized by high sulfate concentrations




- Isolated cropland marsh
- Mix of open water & vegetation(Typha)
- Characterized by lower sulfate concentrations



# LARGE DIFFERENCES IN WATER QUALITY

- **Hogg** higher sulfate, conductivity, DOC, TDN, and ABS 280
- Young higher P & pH



# DIFFERENCES IN G-GFLUXES


Site Year		NEE	FCH4	GHG
		$(gC m^{-2} yr^{-1})$	gC m <sup>-2</sup> yr <sup>-1</sup> ) (g	$gCO_2eq m^{-2} yr^{-1}$
Hogg	2021	-34	1.8	-7
T C	2022	-139	2.5	-349
Young	2021	47	7.9	674
T	2022	-6	4.7	276




# FCH4SPATIAL HETEROGENEITY WITHIN AND BEYOND AFLUX TOWER FOOTPRINT

#### Pairing Flux Maps with Landsat 8

#### Flux Maps




#### Landsat 8






Darian Ng






# COOLING POTENTIAL OF WETLANDS

#### Difference in Diurnal Taero



- Δ T<sub>aero</sub> vary with wetland type
- Cooler T<sub>aero</sub> during daytime and warmer in wetlands
- Cooling upto 4.5 ° C (at noon) and warming upto 2.2 ° C at Young site

#### **Spatial Pattern of LST**





Joyson Ahongshangbam

#### PLANS FOR UPCOMING YEAR

- Install new flux tower in QC
- Continue collaboration with DUC
- Publication of papers on the biogeochemical and biophysical benefits of wetlands

#### POINTS FOR DISCUSSION

- Guidance on lateral flux sampling & water quality data
- Potential collaboration with the QC project
- Integration of data across sites
- Protocols?







## THANK YOU

Sara Knox | sara.knox@mcgill.ca | https://ubc-micromet.github.io/

**ROOM B:** Flux Towers, measurements, and carbon cycling · Christian Von Sperber

**ROOM B:** Flux Towers, measurements, and carbon cycling . Gail Chmura





Related study with Dr. Florin Pendea Lakehead U (Orillia)
Lake Simcoe watershed, ON



winter fluxes critical CH4 emissions when water is frozen over seds not





#### **ROOM C:**

Ecosystem service indicators

## **ROOM C:** Ecosystem service indicators

- . David Aldred
- . Kevin Erratt
- . Owen Salmon
- · Sassan Mohammady
- · Shabnam Majnooni


# **ROOM C:** Ecosystem service indicators


· David Aldred

# Objective 4, Task 2: Develop a desktop- and indicator-based tool for rapid assessment of wetland ecosystem services Irena Creed, David Aldred

In 2013, the Government of Alberta commissioned us to develop a desktop- and indicator-based tool to implement their no-net-loss of wetland function policy → Alberta Wetland Relative Value Evaluation Tool (ABWRET)

Relative wetland values were calculated for over 1 million wetlands in 11 relative wetland value assessment units (RWVAUs) in the province's White Area






#### Main limitations

- (1) variable quality of the wetland database (a mosaic of different inventories captured using different data acquired at different dates and using different specifications)
- (2) varying quality of geospatial indicator data

Due to these limitations, the Government of Alberta reports wetland value categories aggregated to the township level

These are available at:

https://geodiscover.alberta.ca/geoportal/rest/metadata/item/6074617f4be24fcf95a2de56f6f7d28f/html




# **ROOM C:** Ecosystem service indicators

. Kevin Erratt




#### Evaluating urban wetland health: adjusting management strategies

Shift from physical to visual-based assessments, to create a rapid and cost-effective management tool



#### Scaling up: Applying different modelling approaches



Test human vs. machine knowledge to assess to test whether human assistance can enhance machine learning outcomes

#### Modelling Approaches: Hybrid wins (in most instances)

#### **6 Indicators**

#### **10 Indicators**

	Machine	Hybrid	Expert	Machine	Hybrid	Expert
EV						
	0.53	0.78	0.31	0.71	0.6	0.63
14/0						
WQ						
	0.25	0.43	0.4	0.38	0.48	0.44
Н						
	0.29	0.41	0.28	0.42	0.54	0.5
	0.20	<b>5.41</b>	0.20	J.72	0.04	0.0

#### Modelling Approaches: We need human knowledge!

"Human contributions to scientific progress were once deemed to be non-essential and replaceable" in the realm of artificial intelligence Hansen & Quinon (2023) Synthese 201.1-21.

However, we show **hybrid approaches**, which use expert knowledge to "feed" artificial intelligence enhance outcomes.

### **ROOM C:** Ecosystem service indicators

· Owen Salmon

CAAF Objective 4: "Develop robust estimates of the synergies (and conflicts) of wetlands as NBS for carbon storage vs. other benefits"

#### **HYPOTHESIS:**


Rising cyanobacteria dominance is related to % wetland cover in contributing catchments and the associated rise in DOM-Fe supplied to lakes.


#### **MAJOR FINDING:**


Rising cyanobacteria dominance is related to low % wetland cover and high Fe phosphate delivery to lakes, causing an increase in cyanobacteria blooms.


#### Field based study – Summer 2023

- Lakes (n=108) within Alberta,
   Saskatchewan, and Manitoba
- Selected based on % wetland coverage












- 1) The majority of algal biomass is cyanobacteria
- 2) TP drives cyanobacteria, with low N:P ratios (low N environments) regulating cyanobacteria production




- 1) The majority of algal biomass is cyanobacteria
- 2) TP drives cyanobacteria, with low N:P ratios (low N environments) regulating cyanobacteria production
- 3) Aphanizomenon, a N-fixing genera, is dominant. This likely reflects low N conditions



- 1) The majority of algal biomass is cyanobacteria
- 2) TP drives cyanobacteria, with low N:P ratios (low N environments) regulating cyanobacteria production
- 3) Aphanizomenon, a N-fixing genera, is dominant. This likely reflects low N conditions
- 4) No grazing effect




- 1) The majority of algal biomass is cyanobacteria
- 2) TP drives cyanobacteria, with low N:P ratios (low N environments) regulating cyanobacteria production
- 3) Aphanizomenon, a N-fixing genera, is dominant. This likely reflects low N conditions
- 4) No grazing effect
- 5) Terrestrial DOM is fueling cyanobacteria production



- 1) The majority of algal biomass is cyanobacteria
- 2) TP drives cyanobacteria, with low N:P ratios (low N environments) regulating cyanobacteria production
- 3) Aphanizomenon, a N-fixing genera, is dominant. This likely reflects low N conditions
- 4) No grazing effect
- 5) Terrestrial DOM is fueling cyanobacteria production
- 6) Anemic lakes result in high siderophore production





### **ROOM C:** Ecosystem service indicators

· Sassan Mohammady

CAAF Objective 3, Task: 3.2. Estimate The Hydrological Connectivity Of Wetlands To The Watersheds In Which They Are Embedded.

Identifying the role of wetland to river connectivity and wetland loss (or gain) on the headwater lake's phytoplankton biomass (i.e., as an index of water quality )


PhD Candidate: Sassan Mohammady Supervisor: Dr. Irena Creed




#### Hypothesis

Wetland loss within lake catchment will increase phytoplankton biomass in the receiving lake water


#### Study area









# Remote sensing of Chlorophyll-a to create the timeseries of headwater lake's phytoplankton biomass (1984-2023)




## Analyzing the correlation between the timeseries of wetland changes with the timeseries of phytoplankton biomass (i.e., Chl-a)






### **ROOM C:** Ecosystem service indicators

· Shabnam Majnooni





Determine if lakes are losing resilience and transitioning to new stable states, and identify the driving factors.



Alternative stable states within a system











#### **ROOM D:**

Socioeconomic analyses

### **ROOM D:** Socioeconomic analyses

- · John Pattison Williams & Patrick Lloyd Smith & Roy Brouwer
- · Purbasha Mistry

is project was undertaken with the financial support the Government of Canada. Is project a été réalisé avec l'appui financier I gouvernement du Canada.

#### Canada

# Wetland as Natural Solutions: Baselines and projections for Wetlands on Agricultural Land

Patrick Lloyd-Smith and John K. Pattison-Williams
Ashley Klotz, Liam Boldt and .....







#### Overview

**Current Status** 

Plans for Upcoming Season

Challenges

#### **Objectives**

Watershe

• Initial analysis of the Lake Winnipeg watershed, engaging with Ontario/Quebec and the rest of Canada

Wetland Exte Data Collect data on historical, current, and future wetland extent from natural science teams

Drivers of

Identify and describe the main drivers of wetland conversion (e.g. literature review and/or quantitative assessment)

Survey Design

• Design and implement a survey instrument to explore the economic behavioural perspectives of stakeholders towards wetland management and drainage (e.g. wetlands as natural, perceived drivers of loss, risk of leakage).

Economic Mode Development • develop a model that incorporates behaviour to project wetland conversation rates under different scenarios based on economic conditions (e.g. input costs, commodity prices), technology changes (e.g. drainage costs, yields) and policy instruments (e.g. regulations, payment for ecosystem service schemes).

Policy

• Use the developed model and projections to assess the efficiency of various policy instruments


Leakage

Assess the potential role of 'leakage' of wetland associated emissions (i.e., displacement of emissions from avoiding an activity to another location) in the analysis at the sub-provincial level

Administratia

recruitment and initial data collection

### Current Status | Watershed Analysis & Wetland Extent










Relying on David Aldred and team from U of T

### Current Status | Drivers of Conversion: Systematic Review

Stage	Description	Action
1	Google Scholar	Wetlands + Conversion + Agriculture
	Google	Wetlands + Conversion + Drivers
	SCOPUS	Wetlands + Conversion + Urban Expansion
	Science Direct	Wetlands + Conversion + Development
	Web of Science	Wetlands + Conversion + Causes
		Wetlands + Conversion + Forestry
		Wetlands + Conversion + Canada
2	Title and Abstract Review	Document titles and abstracts were reviewed for retention / exclusion
		Criteria: includes explicit wetland loss and discussion of conversion
3	Document Review	Documents were read to explore key themes and relevance.
4	Thematic Analysis	Key messages, lessons and limitations were identified and synthesized for:
		i) context: geographic, grassland type and governance structure (private/public)
		ii) driver of conversion
		iii) ecosystem service explored
		iv) wetland policy structure identified




#### Current Status | Preliminary Search Results





#### Current Status | Systematic Review Thematic Analysis






### Current Status | Economic Model Development

 Improve economic modelling of wetland conservation costs to incorporate uncertainty



 Using precision agriculture data to estimate the effects of wetlands on crop yields within the basin itself as well as in the adjacent buffer areas. Focus is on yield effect differences across soil zones, wet/dry years, wetland sizes, and crop types.

#### Plans for Upcoming Season







### Plans for Upcoming Season | Survey Design



- Reviewing approaches
- Design summer 2023
- Pre-test Fall 2023
- Implement January 2024
- In-person meeting May 2, 2023 in Toronto

### Challenges / Opportunities

Confirming funding and subgrant agreements

Hiring / building team

### **ROOM D:** Socioeconomic analyses

· Purbasha Mistry


## **CAAF Objective 5.4. Support Other National Initiatives For Nature Climate Solutions.**

Wetlands as economically viable nature-based solutions for climate change mitigation


Purbasha Mistry

Supervisors: Drs. Irena Creed and Charles Trick


 Evaluated the value of carbon sequestration in wetlands by calculating the benefit-cost ratio of conserving vs. restoring the wetlands



### Carbon Sequestration Rate of restored and intact wetlands




### Value of carbon incorporates carbon sequestration rate and carbon price of Canada



#### Comparison of costs and benefits to sequester CO<sub>2</sub> in wetlands in Alberta in 2021 \$ CAD

	Present value of costs \$ CAD/ha		Benefit-cost ratio		
					Breakeven cost ( = present
Time periods	Conservation	Restoration	Conservation	Restoration	value of carbon) \$  CAD/ha
20 yrs.	\$2,127	\$25,127	2.59	0.22	\$5, 504
40 yrs.	\$3,305	\$26,305	3.89	0.49	12, 853
50 yrs.	\$3,679	\$26,679	4.42	0.61	\$16,278

The cost of restoring the wetland incorporates restoration cost plus the opportunity cost Used 3% discount rate and 2% inflation



#### **Key Messages:**

- Conservation is more cost-effective than restoration of wetlands
- Incorporating multiple benefits of restored wetlands can help justify the cost

**Your feedback** to improve this economic analysis approach is much appreciated.

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- 3:30 − 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

#### Rose Fuchsia Room, Topic C

#### **Topic:**

**Ecosystem service indicators** 

#### **Objectives:**

CAAF, Objective 4

Great Lakes, Objectives 1, 2, 4.5.

#### **Room leaders:**

David Aldred, James Paterson

#### Purple Room, Topic D

#### Topic:

**Socioeconomic Analyses** 

#### **Objectives:**

CAAF, Objective 5.4.

Prairies: Objective 1 and 2.

Great Lakes: Objectives 3 and 4.

#### **Room leaders:**

John Pattison-Williams, Patrick Lloyd-Smith, Roy Brouwer

#### Rose Fuchsia Room, Topic C

#### **Topic:**

**Ecosystem service indicators** 

#### **CAAF:**

Irena Creed (David Aldred, Kevin Erratt, Sassan Mohammady, Shabnam Majnooni, Owen Salmon, Kelechi Nwokeocha)

James Paterson

Lauren Bortolotti

#### Purple Room, Topic D

#### **Topic:**

**Socioeconomic Analyses** 

#### CAAF:

Irena Creed (Purbasha Mistry\*)

#### **Prairies:**

John Pattison-Williams (Ashley Klotz)

Patrick Lloyd-Smith

#### **Great Lakes:**

Roy Brouwer (Jullian Sone)

Lota Tamini\*

Jie He\*

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- · 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks

- How can we better coordinate activities across projects, objectives, and teams?
- What opportunities can we create for students?
- What synergies within and between projects objectives can we explore?
- Are we on track to achieve our objectives? What needs to be done to address the delays? Are you missing data or information? How can these be provided to you?

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- · 3:30 4:00, From Science to Impact
- $\cdot$  4:00 4:30, Final remarks



#### Task 5.1. CANWIN

Establish an open science Canadian Wetland Information repository to support scientists and decision makers interested in wetlands as nature-based solutions.

Late Start to be mitigated:

• **STEP 1.** Hiring a Business Analyst (May/Jun 2024)
Will interview AAFC, ECCC, Co-PIs to identify requirements for CANWIN (Databases, datasets, normalization, user stories, technical and technological requirements)

• STEP 2. Hiring of Full-Stack application Developer (Expected Nov/Dec 2024)

Challenges: Intellectual property, technological, project afterlife, ...

#### Task 5.2. Wetlands in NIR

Integrate wetlands and their role in OC accumulation and GHG reduction into the national GHG inventory.

- Workshops and collaboration with ECCC AFOLU (Doug MacDonald)
- Wetland mapping for PPR from 1993-2020
- Estimates of wetland conversion rates 1970 to 2020 delivered to ECCC
- Proposal of criteria to classify wetlands as managed/unmanaged according to landproxy approach

# Criteria for classifying wetlands as managed or unmanaged in agricultural landscapes

To report emissions or removals of GHG from lands into the National Inventory Report, they need to be considered MANAGED or caused by DIRECT HUMAN ACTIVITY.

Currently, all wetlands in Canada are considered unmanaged. This impedes the counting of wetlands as nature-based climate change solutions.

Establishing scientific criteria, thresholds, and indicators to distinguish when a wetland has been managed or nor is required to track emissions/removals into NIR.

#### Task 5.3. Wetlands in Holos Model

Develop Approaches To Quantify Agricultural Impacts On Wetland Carbon Storage And Ghg Emissions To Enable Farmers To Calculate Estimates At The Farm Scale So That They Can Make Land Decisions That Are Consistent And Quantifiable At The National Scale (Holos).

• FY 3 Collaboration with AAFC (Roland Kroebel, Sarah Pogue, Shathi Akhter)

Hire postdoctoral fellow to complete literature review, and to develop question structure for wetland inclusion



- Timeline: Start mid-2024, target 2-year duration
- Aim for inclusion in Holos V6 release (2026/2027)

#### Task 5.4. Socioeconomic (Prairies and Great Lakes)

Support Other National Initiatives For Nature Climate Solutions.

Summary of the conclusions from Room D.

- · 8:30 9:00, Welcome
- 9:00 9:30, Introductions
- 9:30 10:30, Breakout room (A, B)
- · 10:30 10:45, Morning Break
- · 10:45 12:00, Breakout rooms (A, B)
- · 12:00 1:00, Lunch break
- · 1:00 2:30, Breakout rooms (C, D)
- · 2:30 2:45, Afternoon Break
- · 2:45 3:30, Breakout rooms (C, D)
- · 3:30 4:00, From Science to Impact
- 4:00 4:30, Final remarks

























Ce projet a été réalisé avec l'appui financier du gouvernement du Canada.

This project was undertaken with the financial support of the Government of Canada.



